

# Microscopic Theory for Emission of Elementary Excitations into He II from a Heated Solid.

Alexander V. Zhukov

*Institute of Physics, Academia Sinica, Nankang, Taipei 11529, Taiwan*

(February 1, 2008)

I develop here the microscopic quantum theory for description of creation of phonons and rotons in superfluid helium by a solid heater. Starting with correct transfer Hamiltonian describing a coupling between the solid and liquid  $^4\text{He}$  the probabilities of transformation of a single phonon in the solid into i) single helium phonon, ii) two helium phonons, and iii) single helium roton are found out. All the obtained expressions account for different polarizations of phonons in the solid. The heat transfer associated with single phonon and single roton channels are calculated. Particularly, the obtained expression for heat flux via the single phonon channel calculated in the framework of present microscopic theory exactly coincides with the well known Khalatnikov formula obtained initially in the framework of acoustic-mismatch theory. The impossibility of direct creation of  $R^{(-)}$  rotons becomes clear in the used framework due to accurate account of the boundary conditions at the solid – liquid helium interface, which is in agreement with recent experimental results.

67.40.-w, 67.40.Pm, 68.08.-p, 05.30.Jp

## I. INTRODUCTION

The elementary excitations in superfluid  $^4\text{He}$  have been the object of extensive study for many years. The phonon–roton type excitation spectrum proposed phenomenologically by Landau<sup>1</sup> allows to account correctly for superfluid properties and thermodynamic behaviour of liquid helium below the  $\lambda$ -transition temperature. The existence of such a spectrum was demonstrated by the numerous direct observations of single excitations from inelastic neutron scattering. In spite of many various attempts to build a completely microscopic theory to derive directly the thermal excitations spectrum, the general picture drawn first by Landau<sup>1</sup> and Feynman and Cohen<sup>2</sup> had not changed essentially during more than four decades.

Since we realized the general picture of elementary thermal excitations in bulk liquid helium, the natural problem appeared: how to describe a heat flow across a boundary between a solid and liquid helium. All earlier investigations of this process were concentrated on studying of a discontinuity in temperature appeared between the two materials<sup>3</sup>, which is commonly known now as Kapitza resistance. The first consistent accurate theory of this effect was developed by Khalatnikov<sup>4</sup>. He built his theory starting with quantizing of elastic waves of different polarizations in the solid. But in general, the main results of Khalatnikov's theory are completely consistent with classical acoustic–mismatch theory due to Little<sup>5</sup>. The in-depth reviews of theoretical and experimental investigations of Kapitza resistance problem are given by Challis<sup>6</sup> and recently by Zinov'eva<sup>7</sup>. In their pure form all these theories show that transfer of phonons from the solid to the liquid helium is impeded first due to a large difference in densities  $\rho_L/\rho_S \ll 1$  ( $\rho_L$  and  $\rho_S$  is the densities of liquid helium and solid, respectively), and second due to the smallness of critical angle  $\sin \theta_c = c/s_t \ll 1$ , where  $c$  is the first sound velocity in helium and  $s_t$  is the velocity of transverse sound in solid. The total internal reflection occurs when the angle of incidence exceeds this angle  $\theta_c$ . Because of this the helium phonon can propagate only into a narrow cone along the direction perpendicular to the interface.

According to the present understanding, solid can inject the phonons in two groups<sup>8</sup>. The first is phonons in the narrow cone perpendicular to the solid heater surface, while the second group is phonons emitted into the entire half space, these are the so-called background phonons. The existence of the first group is understood satisfactory from acoustic–mismatch theories<sup>4–6</sup>. This mechanism was directly confirmed by experiments<sup>9</sup> (see also Ref. 7). However the existence of background phonons, which is evident from experiments on phonon emission from cleaved crystals<sup>9,10</sup>, was not satisfactory explained theoretically until now in spite of many attempts<sup>11</sup>. In fact, all these theories were devoted to various modifications of the acoustic–mismatch theory to account the imperfection of the interface and some additional scattering mechanisms. As a result, only small corrections to the Khalatnikov theory were obtained.

All theoretical works cited above have been based on the classical acoustic theory and there was thus need for a microscopic quantum mechanical description since  $^4\text{He}$  is a quantum liquid rather than a classical elastic fluid. There were several attempts to build a quantum mechanical theory of solid–liquid helium heat transmission based on various simplified models<sup>12</sup>, but the first really essential contribution to the subject has been done by Sheard, Bowley, and Toombs<sup>13</sup> (hereinafter is referred as SBT). In fact I present here the modification of this theory.

The SBT theory is based on the description of coupling between the solid and the liquid  $^4\text{He}$  via a transfer Hamiltonian in analogy with the theory of electron tunneling through a metal-insulator-metal junction. The main insight of Ref. 13 consist in the expression of tunnel Hamiltonian through the displacement of the solid at the interface and the stress tensor of liquid helium.

The main shortcoming of SBT theory is the clear internal asymmetry of their transfer Hamiltonian with respect to solid and He II. Namely, as it follows from their theory, the solid acts on the helium, but there is no return action. As a result the transfer coefficient for the single phonon channel (single solid phonon – single helium phonon) differs from the result obtained by Khalatnikov<sup>4</sup>, which is correct for this process without any doubt. The authors of Ref. 13 have also roughly estimated the transfer coefficients for single solid phonon – two helium phonons (two phonons process) and single solid phonon – single roton processes. As it will be shown below the more accurate computation of these coefficients gives rather different results.

Calculation of probability for creation of rotons by a solid heater is called for by recent experiments of Exeter group<sup>14</sup>, which obtained quite a number of very interesting results. Particularly, they concluded that creation of the  $R^{(+)}$  rotons (rotons with wave vector greater than corresponding to roton minimum  $k_0 \approx 1.92\text{\AA}^{-1}$ ) does not essentially depend on the particular geometry of the heater, while the  $R^{(-)}$  rotons (rotons with wave vector smaller than corresponding to roton minimum  $k_0 \approx 1.92\text{\AA}^{-1}$ ) cannot be created by a simple flat heater at all. This observation still requires the proper explanation.

In this paper I propose a microscopic quantum theory for creation of thermal excitations in superfluid  $^4\text{He}$  by a heated solid. The starting point of the developed theory is the construction of more correct transfer Hamiltonian to exploit further the main idea of SBT theory<sup>3</sup>. To test the developed theory I first consider the heat flow via the single phonon channel with account of all phonon polarizations in a solid. The obtained expression for a heat flux through the interface associated with this process coincides exactly with the well known Khalatnikov result<sup>4</sup>. This direct support of the formalism allowed me to consider also the single roton emission process as well as to write down the probabilities for emission of a phonon pair by a single phonon in the solid. Particularly, I argue that emission of  $R^{(-)}$  rotons is forbidden by the boundary conditions at the solid – He II interface, which is in agreement with experimental observations<sup>14</sup>.

The paper is constructed as follows. The transfer Hamiltonian corresponding to solid – liquid helium interface is constructed in Section II. In Section III I write down the tunnel Hamiltonians for the particular single excitation and double excitations processes. The well-known Khalatnikov formula for a transfer of single solid phonon into single phonon in helium is obtained from the general formalism in Section IV. Finally in Section V I study the creation of rotons from a heated solid via the single excitation channel. The outlines and conclusions are given in Section VI.

## II. SOLID–LIQUID COUPLING HAMILTONIAN

The whole system under consideration consists of the solid and the superfluid helium with a common boundary between them. To study the excitations transmission processes through the interface we must pick up the part of total Hamiltonian, which corresponds to the interface itself. Just this interface Hamiltonian represents the coupling between two media.

Let  $\hat{\psi}^\dagger(\mathbf{r})$  and  $\hat{\psi}(\mathbf{r})$  be the field operators (creation and annihilation operators in Schrödinger representation) in helium. Here  $\mathbf{r}$  is the coordinate of Lagrange point in superfluid helium. Suppose that thermal perturbation lead to the displacement of some atom in solid with equilibrium coordinate  $\mathbf{R}_n$  by vector  $\mathbf{u}_n$ . In the frame of pair interaction approximation we can write the deviation of a total Hamiltonian from the equilibrium one in the following form:

$$\hat{\mathcal{H}}_{\text{int}} = \int_{\Omega_L} \hat{\psi}^\dagger(\mathbf{r}) \sum_n \hat{U}_{\text{int}}(|\mathbf{R}_n + \mathbf{u}_n - \mathbf{r} - \delta\mathbf{r}|) \hat{\psi}(\mathbf{r}) d^3r, \quad (1)$$

where  $\Omega_L$  is the volume of the helium,  $\hat{U}_{\text{int}}$  is the potential of interaction between a physical point of helium with a volume element in solid. This potential behaves like a single-particle operator when act on the creation and annihilation operators in helium. The quantity  $\delta\mathbf{r}$  in eq. (1) stands for the physical point displacement in helium due to the perturbation  $\mathbf{u}_n$  in solid. It is connected with the density fluctuations in helium by the relation

$$\delta n(\mathbf{r}) = n(\mathbf{r}) - n_0 = \frac{\partial n}{\partial \mathbf{r}} \delta\mathbf{r}. \quad (2)$$

Let us assume the interface to be a plane perpendicular to  $z$ th direction as is shown in Fig. 1. Supposing the displacements  $\mathbf{u}$  and  $\delta\mathbf{r}$  to be small, we can rewrite expression (1) up to the first order in these perturbations as

$$\hat{\mathcal{H}}_{\text{int}} = \int_{\Omega_L} d^3r n(\mathbf{r}) U_0(\mathbf{r}) + \int_{\Omega_L} d^3r u_z(x, y, 0) f_z^{(SL)}(\mathbf{r}) - \int_{\Omega_L} d^3r \delta r_z(x, y, 0) f_z^{(LS)}(\mathbf{r}), \quad (3)$$

where

$$\mathbf{f}^{(SL)}(\mathbf{r}) = \hat{\psi}^\dagger(\mathbf{r}) \sum_n \nabla_{\mathbf{R}_n} \hat{U}_{\text{int}}(\mathbf{R}_n - \mathbf{r}) \hat{\psi}(\mathbf{r}), \quad (4)$$

$$\mathbf{f}^{(LS)}(\mathbf{r}) = \hat{\psi}^\dagger(\mathbf{r}) \nabla_{\mathbf{r}} \sum_n \hat{U}_{\text{int}}(\mathbf{R}_n - \mathbf{r}) \hat{\psi}(\mathbf{r}) \quad (5)$$

are the forces per unit volume of liquid helium and solid, respectively, and

$$U_0(\mathbf{r}) = \sum_n \hat{U}_{\text{int}}(\mathbf{R}_n - \mathbf{r}).$$

In the expression (3) we take the quantities  $u_z(x, y, 0)$  and  $\delta r_z(x, y, 0)$  at the plane  $z = Z = 0$  because of assumed short-range interaction. The first term in eq. (3) corresponds to a static interaction and does not depend on the local spatial perturbations. Thus, it naturally should be included to the equilibrium Hamiltonian. To transform remaining two terms we have to account correctly the boundary conditions at the interface. First, require the normal components of displacements in solid and liquid helium at the surface to be equal

$$u_z(x, y, 0) = \delta r_z(x, y, 0). \quad (6)$$

Second, the total work done must be zero. This requirement can be expressed mathematically as

$$\int dz f_z^{(SL)}(x, y, z) + \int dZ f_z^{(LS)}(X, Y, Z) = 0. \quad (7)$$

Using the relations (6) and (7) we can express the interaction Hamiltonian (3) in the following simple form

$$\hat{\mathcal{H}}_{\text{int}} = 2 \int_{\Omega_L} d^3r u_z(x, y, 0) f_z^{(SL)}(\mathbf{r}). \quad (8)$$

Note, the interaction Hamiltonian (8) obtained here differs from SBT Hamiltonian<sup>13</sup> by the factor 2. This difference follows from incorrect account of mutual influence of solid and liquid helium by the authors of Ref. 13. They have accounted only the perturbation in the solid, which leads to an incident excitation. But as it can be seen from the expression (1) above, the account of corresponding perturbation in helium leads to a return response, which doubles the interaction Hamiltonian in view of the boundary conditions (6) and (7).

Using the relation

$$\nabla_{\mathbf{R}_n} \hat{U}_{\text{int}}(\mathbf{R}_n - \mathbf{r}) = -\nabla_{\mathbf{r}} \hat{U}_{\text{int}}(\mathbf{r} - \mathbf{R}_n)$$

and the expression (4) for  $\mathbf{f}^{(SL)}$  we can get the following relation

$$f_z^{(SL)}(x, y, z) \approx -n(\mathbf{r}) \frac{dU_0}{dz}, \quad (9)$$

where  $n(\mathbf{r})$  is the number density of helium atoms. Now we come to the first essential approximation. Let us choose  $U_0(z)$  in the form of repulsive potential step at the solid-liquid interface

$$U_0(z) = \begin{cases} V_0 & (z < 0) \\ 0 & (z > 0) \end{cases}, \quad (10)$$

where  $V_0 = \text{const} > 0$ . Below I will use the method of elimination of the force (9) from Hamiltonian (8) proposed by SBT<sup>13</sup>. Due to their insight the height  $V_0$  of the step (10) is not crucial. Note, the combination  $-n(\mathbf{r})(dU_0/dz)$  is the external force density exerted on the liquid helium by the solid. The hydrodynamic equation of motion looks

$$\frac{\partial j_k}{\partial t} + \frac{\partial \pi_{kl}}{\partial x_l} = f_k^{(SL)}, \quad (11)$$

where  $\mathbf{j}$  is the momentum density of liquid helium,  $\pi_{kl}$  is the fluid stress tensor. On the other hand the quantum mechanical equation for evolution of operator  $\mathbf{j}$  (Heisenberg equation) gives

$$\frac{i}{\hbar} [j_z, \hat{\mathcal{H}}_L] = n(\mathbf{r}) \frac{dU_0}{dz} + \frac{\partial \pi_{zl}}{\partial x_l}, \quad (12)$$

where  $\hat{\mathcal{H}}_L$  is the Hamiltonian of liquid helium. Following Ref. 13 let us integrate equation (12) in the interval  $z \in \{-\varepsilon, +\varepsilon\}$ , where  $\varepsilon$  is of the order of distance over which an average helium density decays from its bulk value to zero inside the potential step. In fact, I will put finally  $\varepsilon \rightarrow 0$ , which is consistent with our choice of the potential (10). Since the momentum density  $\mathbf{j}$  remains finite the integral over the left hand side of eq. (12) behaves like  $o(z)$  and therefore can be neglected. The derivatives of all components of vector  $\pi_{zl}$  except  $\partial \pi_{zz}/\partial z$  are finite as well. So, we get

$$-\int_{-\varepsilon}^{+\varepsilon} n(\mathbf{r}) \frac{dU_0}{dz} dz = \pi_{zz} \Big|_{z=\varepsilon} - \pi_{zz} \Big|_{z=-\varepsilon} = \pi_{zz}(x, y, 0). \quad (13)$$

Finally, using the formulae (8), (9), and (13) we can write the interaction Hamiltonian in the form

$$\hat{\mathcal{H}}_{\text{int}} = 2 \int_{\Sigma} u_z(x, y, 0) \pi_{zz}(x, y, 0) dx dy, \quad (14)$$

where  $\Sigma$  is the area of solid-liquid interface. Formula (14) represents the interaction Hamiltonian, expressed through the solid displacement at the surface and diagonal component of liquid stress tensor. Now it should be represented in terms of collective variables, which allow to carry out second quantization in the simplest way. It is naturally to present  $\pi_{zz}$  as a sum of kinetic and potential parts as follows

$$\pi_{zz} = \pi_{zz}^{(\text{kin})} + \pi_{zz}^{(\text{pot})}. \quad (15)$$

The explicit forms of  $\pi_{zz}^{(\text{kin})}$  and  $\pi_{zz}^{(\text{pot})}$  are derived in Ref. 13 using the semi-microscopic theory of superfluids developed by Sunakawa *et al.*<sup>15</sup>. Thus, in our notations the kinetic part of (15) takes the form

$$\pi_{zz}^{(\text{kin})} = mn v_z^2 + \frac{\hbar^2}{2m} \left( \frac{\partial n}{\partial z} n^{-1} \frac{\partial n}{\partial z} - \frac{\partial^2 n}{\partial z^2} \right) - \sum_{\mathbf{k}} \frac{\hbar^2 k_z^2}{2m\Omega_L}, \quad (16)$$

where  $m$  is the mass of helium atom,  $n$  is the number density operator, which stands for the first collective variable,  $v_z$  is  $z$ th component of the velocity operator, which stands for the second collective variable,  $\mathbf{k}$  is the wave vector of excitation in liquid helium.

The potential part of (15) can be obtained from the equation of motion, which leads to the relation

$$\sum_{\mathbf{k}} \frac{\partial \pi_{zz}^{(\text{pot})}}{\partial x_{\mathbf{k}}} = \hat{\psi}^\dagger(\mathbf{r}) \left\{ \int d^3 r' \hat{\psi}^\dagger(\mathbf{r}') V(\mathbf{r} - \mathbf{r}') \hat{\psi}(\mathbf{r}') \right\} \hat{\psi}(\mathbf{r}), \quad (17)$$

where  $V(\mathbf{r} - \mathbf{r}')$  is the inter-atomic potential in helium. In fact the equation (17) implies the pair interaction, which strictly speaking is inappropriate for the liquid helium. We however will eliminate it from all final relations by the use of semi-phenomenological notations associated with superfluid helium. The detailed derivation of  $\pi_{zz}^{(\text{pot})}$  from eq. (17) can be found in Ref. 13.

In terms of Fourier transforms defined by  $\mathbf{v} = \sum_{\mathbf{k}} \mathbf{v}_{\mathbf{k}} \exp(i\mathbf{k}\mathbf{r})$  and  $n = \sum_{\mathbf{k}} n_{\mathbf{k}} \exp(-i\mathbf{k}\mathbf{r})$  the kinetic and potential parts of  $z$ th diagonal element of the stress tensor up to the second order in collective variables look

$$\begin{aligned} \pi_{zz}^{(\text{kin})} = \sum_{\mathbf{k}, \mathbf{k}'} & \left\{ \rho_L v_{-\mathbf{k}z} v_{-\mathbf{k}'z} - \frac{\hbar^2 k_z k'_z}{4\rho_L} n_{\mathbf{k}z} n_{\mathbf{k}'z} \right\} e^{-i(\mathbf{k}+\mathbf{k}')\mathbf{r}} \\ & + \frac{\hbar^2}{4m} \sum_{\mathbf{k}} k_z^2 n_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}} - \sum_{\mathbf{k}} \frac{\hbar^2 k_z^2}{2m\Omega_L}, \end{aligned} \quad (18)$$

$$\pi_{zz}^{(\text{pot})} = \sum_{\substack{\mathbf{k}, \mathbf{p} \\ \mathbf{p} \neq 0}} n_{\mathbf{p}-\mathbf{k}} n_{\mathbf{k}} V(\mathbf{k}) \frac{k_z}{p_z} e^{-i\mathbf{p}\mathbf{r}} \quad (19)$$

where  $V(\mathbf{k})$  is the Fourier transform of interatomic interaction potential in helium, which will be expressed below in terms of excitation spectrum of superfluid helium. Above we have used only two essential approximations: the step shape of potential (10) and the pair interaction between helium atoms which results in the expression (19). The first approximation can be naturally justified in view of smallness of boundary inhomogeneity domain, which is of the order of interatomic distances<sup>16</sup>. To avoid the explicit use of the second approximation I will act as follows. The most general approach to the problem of low lying energy levels in superfluids lead to the following form of excitation spectrum<sup>17</sup>

$$\hbar\omega_{\mathbf{k}} = \frac{\hbar^2 k^2}{2m\Lambda_{\mathbf{k}}}, \quad (20)$$

where  $\hbar\omega_{\mathbf{k}}$  is the energy of excitation with momentum  $\hbar\mathbf{k}$ ,  $\Lambda_{\mathbf{k}}$  is the structure factor of superfluid helium. Within the framework of microscopic theory<sup>15</sup> using the approximation of pair interaction this structure factor takes the form

$$\Lambda_{\mathbf{k}} = \left\{ 1 + 4 \frac{\rho_L V(\mathbf{k})}{\hbar^2 k^2} \right\}^{-1/2}. \quad (21)$$

This particularly means that if we know the correct excitation spectrum of superfluid helium, say from experiment, then the quantity  $V(\mathbf{k})$  can be unambiguously expressed through the parameters of this spectrum. Namely

$$V(\mathbf{k}) = \frac{1}{\rho_L} \left( \frac{m^2 \omega_{\mathbf{k}}^2}{k^2} - \frac{1}{4} \hbar^2 k^2 \right), \quad (22)$$

where the analytic form of  $\omega_{\mathbf{k}}$  depends on particular range of wave vectors. In this manner the problem becomes self-consistent.

The Fourier transforms of collective variables can now be written in the second quantized form<sup>18</sup>

$$n_{-\mathbf{k}} = -i \left( \frac{\rho_L \Lambda_{\mathbf{k}}}{m \Omega_L} \right)^{1/2} (\hat{a}_{\mathbf{k}} - \hat{a}_{-\mathbf{k}}^\dagger) = -i \frac{\hbar k}{m} \left( \frac{\rho_L}{2 \hbar \omega_{\mathbf{k}} \Omega_L} \right)^{1/2} (\hat{a}_{\mathbf{k}} - \hat{a}_{-\mathbf{k}}^\dagger). \quad (23)$$

$$\mathbf{v}_{\mathbf{k}} = -\frac{i}{2} \frac{\hbar \mathbf{k}}{\sqrt{m \rho_L \Omega_L \Lambda_{\mathbf{k}}}} (\hat{a}_{\mathbf{k}} + \hat{a}_{-\mathbf{k}}^\dagger) = -i \left( \frac{\hbar \omega_{\mathbf{k}}}{2 \rho_L \Omega_L} \right)^{1/2} \frac{\mathbf{k}}{k} (\hat{a}_{\mathbf{k}} + \hat{a}_{-\mathbf{k}}^\dagger). \quad (24)$$

where  $\hat{a}_{\mathbf{k}}^\dagger$  and  $\hat{a}_{\mathbf{k}}$  are the creation and annihilation operators of elementary excitation in liquid helium with momentum  $\hbar\mathbf{k}$  and energy  $\omega_{\mathbf{k}}$ .

Now we must write down the  $z$ th component of lattice displacement in the solid in terms of creation  $\hat{b}_{\mathbf{q}_\alpha}^\dagger$  and annihilation  $\hat{b}_{\mathbf{q}_\alpha}$  operators for solid phonons. Here  $\mathbf{q}_\alpha$  is the wave vector of solid phonon with  $\alpha$ th polarization. Taking into account the longitudinal and transverse polarizations of bulk phonons and surface phonons we can write

$$u_z(\mathbf{R}) = \left( \frac{\hbar}{2 \rho_S \Omega_S} \right)^{1/2} \left\{ \sum_{\mathbf{q}_l} \frac{B_l(\theta)}{\omega_{\mathbf{q}_l}^{1/2}} (\hat{b}_{\mathbf{q}_l} - \hat{b}_{-\mathbf{q}_l}^\dagger) e^{i\mathbf{q}_l \mathbf{R}} + \sum_{\mathbf{q}_t} \frac{B_t(\theta)}{\omega_{\mathbf{q}_t}^{1/2}} (\hat{b}_{\mathbf{q}_t} - \hat{b}_{-\mathbf{q}_t}^\dagger) e^{i\mathbf{q}_t \mathbf{R}} \right. \\ \left. + \left( \frac{\Omega_S}{\Sigma} \right)^{1/2} \sum_{\mathbf{q}_s} \frac{B_s(q_s)}{\omega_{\mathbf{q}_s}^{1/2}} (\hat{b}_{\mathbf{q}_s} - \hat{b}_{-\mathbf{q}_s}^\dagger) e^{i\mathbf{q}_s \mathbf{r}_\parallel} \right\}, \quad (25)$$

where  $\rho_S$  and  $\Omega_S$  are the mass density and volume of the solid, respectively,  $\omega_{\mathbf{q}_\alpha}$  is the frequency of  $\alpha$ th phonon mode,  $\mathbf{q}_s$  is the two-dimensional wave vector of surface phonon,  $\mathbf{r}_\parallel = \mathbf{R}_\parallel$  is the parallel component of vector  $\mathbf{r}$  (or  $\mathbf{R}$ ) at  $z = Z = 0$ . The angular amplitudes  $B_\alpha(\theta)$  can be determined as<sup>4</sup>

$$B_l(\theta) = \frac{2s_l^2 \cos \theta_0 \cos 2\theta_t}{s_t^2 \sin 2\theta_t \sin 2\theta_0 + s_l^2 \cos^2 2\theta_t}, \quad (26)$$

$$B_t(\theta) = \frac{2s_t^2 \cos \theta_0 \sin 2\theta_l}{s_t^2 \sin 2\theta_l \sin 2\theta_0 + s_l^2 \cos^2 2\theta_0}, \quad (27)$$

$$B_s(q_s) = \frac{q_s^2 - \kappa_t^2}{2q_s} g^{-1/2}, \quad (28)$$

$$g = \frac{q_s^2 + \kappa_t^2}{2\kappa_t} + \frac{q_s^2 + \kappa_t^2}{2\kappa_l} \frac{\kappa_t}{\kappa_l} - \frac{q_s^2 + \kappa_t^2}{\kappa_l} \quad (29)$$

$$\kappa_\alpha = \omega_{\mathbf{q}_\alpha} \sqrt{c^{-2} - s_\alpha^{-2}}, \quad (\alpha = l, t). \quad (30)$$

Here  $c$  is the velocity of first sound in superfluid helium,  $s_l$  and  $s_t$  are the velocities of longitudinal and transverse sounds in the solid,  $\theta_0$  and  $\theta_\alpha$  are the incidence and reflection angles.

In fact, the formulae (14), (18)–(21), and (25) give us the sought transfer Hamiltonian in second quantized form, which describes the transmission of elementary excitations from the solid to superfluid helium.

Note, the resulting Hamiltonian contains the terms associated with creation or annihilation of single excitation as well as pair of excitations in helium, while it is only linear in creation and annihilation operators of phonons in the solid. This asymmetry is of course the shortcoming of our method. It can be avoided if we take into account the third order anharmonicities in the solid. However I am not concerned with this problem here because the main goal of the present paper is to describe the transmission of excitations from *heated solid* to the liquid helium at the temperatures closed to zero, which is consistent with real experimental situation<sup>14</sup>. Under such conditions the contribution of single solid phonon annihilation is expected to dominate.

### III. TRANSFER HAMILTONIANS FOR PARTICULAR PROCESSES

In this Section I write down the Hamiltonians for particular transfer processes: solid phonon  $\Leftrightarrow$  single liquid phonon (single phonon emission), solid phonon  $\Leftrightarrow$  two liquid phonons (two phonons emission), solid phonon  $\Leftrightarrow$  single liquid roton (single roton emission).

#### A. Single phonon emission

To obtain the transfer Hamiltonian corresponding to the tunneling of a single phonon from the solid to the liquid helium and *vice versa* (*i.e.* to the processes  $\mathbf{q} \leftrightarrow \mathbf{k}$ ) we must retain in the expressions (18) and (19) only liner terms with respect to helium collective variable. In this case we get

$$\pi_{zz}^{(1\text{ph})} = \sum_{\mathbf{k}} \left( \frac{\hbar^2 k_z^2}{4m} + \frac{NV(\mathbf{k})}{\Omega_L} \right) n_{-\mathbf{k}} e^{i\mathbf{k}\mathbf{r}}, \quad (31)$$

where I have used  $n_0 = N/\Omega_L$ ,  $N$  is the total number of helium atoms. In the case of helium phonons  $\omega_{\mathbf{k}} = c\mathbf{k}$ . Thus the relation (22) yields

$$V(\mathbf{k}) = \frac{\Omega_L}{N} \left( mc^2 - \frac{\hbar^2 k^2}{4m} \right) \approx \frac{\Omega_L}{N} mc^2, \quad (32)$$

where I have neglected the term quadratic in wave vector. The same approximation can be done in the first term of eq. (31). In fact, without these approximations one can see that after substitution of eq. (32) into eq. (31) both small terms give even smaller one  $\sim (k^2 - k_z^2)$ , because  $k \sim k_z$  in the process of interest. So, using the relations (32) and (31) we come to

$$\pi_{zz}^{(1\text{ph})} = mc^2 \sum_{\mathbf{k}} n_{-\mathbf{k}} e^{i\mathbf{k}\mathbf{r}}. \quad (33)$$

Substituting eqs. (25) and (33) into eq. (14) and integrating over the interface area  $\Sigma$  we obtain the transfer Hamiltonian for the single phonon channel in the following form

$$\begin{aligned}
\hat{\mathcal{H}}_{\text{int}}^{(1\text{ph})} = i\Sigma c\hbar \left( \frac{\rho_L}{\rho_S \Omega_S \Omega_L} \right)^{1/2} & \left\{ \sum_{\mathbf{k}, \mathbf{q}_l} B_l(\theta) \left( \frac{\omega_{\mathbf{k}}}{\omega_{\mathbf{q}_l}} \right)^{1/2} \left( \hat{a}_{\mathbf{k}} \hat{b}_{\mathbf{q}_l}^\dagger - \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{q}_l} \right) \delta_{\mathbf{k} \parallel, \mathbf{q}_l \parallel} \right. \\
& + \sum_{\mathbf{k}, \mathbf{q}_t} B_t(\theta) \left( \frac{\omega_{\mathbf{k}}}{\omega_{\mathbf{q}_t}} \right)^{1/2} \left( \hat{a}_{\mathbf{k}} \hat{b}_{\mathbf{q}_t}^\dagger - \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{q}_t} \right) \delta_{\mathbf{k} \parallel, \mathbf{q}_t \parallel} \\
& \left. + \left( \frac{\Omega_S}{\Sigma} \right)^{1/2} \sum_{\mathbf{k}, \mathbf{q}_s} B_s(q_s) \left( \frac{\omega_{\mathbf{k}}}{\omega_{\mathbf{q}_s}} \right)^{1/2} \left( \hat{a}_{\mathbf{k}} \hat{b}_{\mathbf{q}_s}^\dagger - \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{q}_s} \right) \delta_{\mathbf{k} \parallel, \mathbf{q}_s \parallel} \right\}. \tag{34}
\end{aligned}$$

This formula will be used in Section IV for deriving the Khalatnikov formula for heat flow via the single phonon channel.

## B. Two phonons emission

The obtained formulae (14), (18), and (19) allow us to study also the processes of emission of two helium phonons from single phonon in the solid. For this purpose let us retain in eqs. (18) and (19) only the terms quadratic in collective variables of liquid helium. This leads to the following expression for stress tensor

$$\pi_{zz}^{(2\text{ph})}(\mathbf{r}) = \sum_{\mathbf{k}, \mathbf{k}'} \left\{ \rho_L v_{\mathbf{k}z} v_{\mathbf{k}'z} + \frac{V(\mathbf{k})k_z}{k_z + k'_z} n_{-\mathbf{k}} n_{-\mathbf{k}'} \right\} e^{i(\mathbf{k}+\mathbf{k}')\mathbf{r}}, \tag{35}$$

which gives the transfer Hamiltonian for the two phonons processes under consideration in the form

$$\begin{aligned}
\hat{\mathcal{H}}_{\text{int}}^{(2\text{ph})} = \frac{2\hbar\Sigma}{\Omega_L} \left( \frac{\hbar}{2\rho_S \Omega_S} \right)^{1/2} & \left\{ \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}_l} B_l(\theta) \left( \frac{\omega_{\mathbf{k}} \omega_{\mathbf{k}'}}{\omega_{\mathbf{q}_l}} \right)^{1/2} \Xi(\mathbf{k}, \mathbf{k}', \mathbf{q}_l) \right. \\
& + \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}_t} B_t(\theta) \left( \frac{\omega_{\mathbf{k}} \omega_{\mathbf{k}'}}{\omega_{\mathbf{q}_t}} \right)^{1/2} \Xi(\mathbf{k}, \mathbf{k}', \mathbf{q}_t) \\
& \left. + \left( \frac{\Omega_S}{\Sigma} \right)^{1/2} \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}_s} B_s(q_s) \left( \frac{\omega_{\mathbf{k}} \omega_{\mathbf{k}'}}{\omega_{\mathbf{q}_s}} \right)^{1/2} \Xi(\mathbf{k}, \mathbf{k}', \mathbf{q}_s) \right\}. \tag{36}
\end{aligned}$$

Here I have introduced the functions

$$\Xi(\mathbf{k}, \mathbf{k}', \mathbf{q}_\alpha) = \mathcal{J}_1(\mathbf{q}_\alpha, \mathbf{k}|\mathbf{k}') \delta_{\mathbf{q}_\alpha \parallel + \mathbf{k} \parallel, \mathbf{k}' \parallel} - \mathcal{J}_2(\mathbf{q}_\alpha | \mathbf{k}, \mathbf{k}') \delta_{\mathbf{q}_\alpha \parallel, \mathbf{k} \parallel + \mathbf{k}' \parallel}, \tag{37}$$

where

$$\mathcal{J}_1(\mathbf{q}_\alpha, \mathbf{k}|\mathbf{k}') = \left( \frac{1}{2} + \frac{k_z k'_z}{k k'} \right) \left( \hat{b}_{\mathbf{q}_\alpha} \hat{a}_{\mathbf{k}} \hat{a}_{\mathbf{k}'}^\dagger + \hat{b}_{\mathbf{q}_\alpha}^\dagger \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}'} \right), \tag{38}$$

$$\mathcal{J}_2(\mathbf{q}_\alpha | \mathbf{k}, \mathbf{k}') = \frac{1}{2} \left( \frac{k_z k'_z}{k k'} + \frac{k_z}{k_z + k'_z} \right) \left( \hat{b}_{\mathbf{q}_\alpha} \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}'}^\dagger + \hat{b}_{\mathbf{q}_\alpha}^\dagger \hat{a}_{\mathbf{k}} \hat{a}_{\mathbf{k}'} \right). \tag{39}$$

Formulae (36) – (39) provide a possibility to study the energy transfer through the solid – liquid helium interface without conservation of number of quasiparticles. In this case there is no strong restriction on the angles for created phonons. Therefore such processes can lead to the isotropic background, which has been observed in experiments on phonon emission from cleaved crystals<sup>9,10</sup>. This problem will be considered in detail in another paper.

## C. Single roton emission

To obtain the tunnel Hamiltonian, which corresponds to creation of a single roton in He II by a solid phonon, we can again use the relation (31) with account of eq. (22). In this case we however cannot neglect the terms quadratic in wave vectors. Note, the frequency  $\omega_{\mathbf{k}}$  in (31) represents now the roton spectrum, whose analytic form depends on

the chosen concrete model. Here I do not distinguish the  $R^{(-)}$  and  $R^{(+)}$  rotons. This problem will be considered in detail in Section V.

Substituting eq. (22) into eq. (31) we get

$$\pi_{zz}^{(1R)}(\mathbf{r}) = \sum_{\mathbf{k}} \left\{ \frac{m\omega_{\mathbf{k}}^2}{k^2} + \frac{\hbar^2(k_z^2 - k^2)}{4m} \right\} n_{-\mathbf{k}} e^{i\mathbf{k}\mathbf{r}}. \quad (40)$$

Finally the corresponding transfer Hamiltonian takes the form

$$\begin{aligned} \hat{\mathcal{H}}_{\text{int}}^{(1R)} = & \frac{i\Sigma\hbar}{2m} \left( \frac{\rho_L}{\rho_S\Omega_S\Omega_L} \right)^{1/2} \left\{ \sum_{\mathbf{k}, \mathbf{q}_l} \frac{B_l(\theta)\mathcal{F}(\mathbf{k})k}{(\omega_{\mathbf{k}}\omega_{\mathbf{q}_l})^{1/2}} \left( \hat{a}_{\mathbf{k}}\hat{b}_{\mathbf{q}_l}^\dagger - \hat{a}_{\mathbf{k}}^\dagger\hat{b}_{\mathbf{q}_l} \right) \delta_{\mathbf{k}_{\parallel}, \mathbf{q}_{l\parallel}} \right. \\ & + \sum_{\mathbf{k}, \mathbf{q}_t} \frac{B_t(\theta)\mathcal{F}(\mathbf{k})k}{(\omega_{\mathbf{k}}\omega_{\mathbf{q}_t})^{1/2}} \left( \hat{a}_{\mathbf{k}}\hat{b}_{\mathbf{q}_t}^\dagger - \hat{a}_{\mathbf{k}}^\dagger\hat{b}_{\mathbf{q}_t} \right) \delta_{\mathbf{k}_{\parallel}, \mathbf{q}_{t\parallel}} \\ & \left. + \left( \frac{\Omega_S}{\Sigma} \right)^{1/2} \sum_{\mathbf{k}, \mathbf{q}_s} \frac{B_s(q_s)\mathcal{F}(\mathbf{k})k}{(\omega_{\mathbf{k}}\omega_{\mathbf{q}_s})^{1/2}} \left( \hat{a}_{\mathbf{k}}\hat{b}_{\mathbf{q}_s}^\dagger - \hat{a}_{\mathbf{k}}^\dagger\hat{b}_{\mathbf{q}_s} \right) \delta_{\mathbf{k}_{\parallel}, \mathbf{q}_{s\parallel}} \right\}, \end{aligned} \quad (41)$$

where

$$\mathcal{F}(\mathbf{k}) = \left\{ \frac{m\omega_{\mathbf{k}}^2}{k^2} + \frac{\hbar^2(k_z^2 - k^2)}{4m} \right\}. \quad (42)$$

In Section V I will use the Hamiltonian (41) for derivation of energy flux through the interface via the single roton channel.

As the curtain fell of this section it should be noted that using the general formulae (18) and (19) we could study also various inelastic processes with rotons, *e.g.* solid phonon  $\Leftrightarrow$  two rotons, solid phonon  $\Leftrightarrow$  roton + phonon in helium, and scattering of roton on the surface with creation of a phonon in the solid.

#### IV. SINGLE PHONON EMISSION. DERIVATION OF THE KHALATNIKOV FORMULA

Any new theory claiming to describe the emission of elementary excitations from the solid to the superfluid helium should first be tested on the basic result obtained by Khalatnikov for a heat flow via the single phonon channel<sup>4</sup>. Really, this latter result was derived within appropriate framework, and is in excellent agreement with the experimental observations<sup>9,7</sup>.

To calculate the energy flux through the solid-liquid interface via single phonon channel we must first calculate the probability density function defined by the usual relation

$$W^{(1\text{ph})} = \frac{2\pi}{\Sigma\hbar^2} \left| \mathcal{M}_{\mathbf{q} \leftrightarrow \mathbf{k}}^{(1\text{ph})} \right|^2 \delta(\omega_{\mathbf{q}} - \omega_{\mathbf{k}}), \quad (43)$$

where  $\mathcal{M}_{\mathbf{q} \leftrightarrow \mathbf{k}}^{(1\text{ph})} = \langle \mathbf{q} | \hat{\mathcal{H}}_{\text{int}}^{(1\text{ph})} | \mathbf{k} \rangle$  is the corresponding matrix element of the transfer Hamiltonian given by eq. (34). Substituting (34) into (43) we easily get

$$W^{(1\text{ph})} = (2\pi)^3 \left\{ \mathcal{A}_l^{(1\text{ph})} \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{l\parallel}) + \mathcal{A}_t^{(1\text{ph})} \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{t\parallel}) + \mathcal{A}_s^{(1\text{ph})} \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{s\parallel}) \right\} \delta(\omega_{\mathbf{q}} - \omega_{\mathbf{k}}), \quad (44)$$

where the coefficients  $\mathcal{A}_{\alpha}$  are defined by the relations

$$\mathcal{A}_l^{(1\text{ph})} = \frac{\omega_{\mathbf{k}}}{\omega_{\mathbf{q}_l}} \frac{\rho_L c^2}{\rho_S \Omega_L \Omega_S} B_l^2(\theta), \quad \mathcal{A}_t^{(1\text{ph})} = \frac{\omega_{\mathbf{k}}}{\omega_{\mathbf{q}_t}} \frac{\rho_L c^2}{\rho_S \Omega_L \Omega_S} B_t^2(\theta), \quad \mathcal{A}_s^{(1\text{ph})} = \frac{\omega_{\mathbf{k}}}{\omega_{\mathbf{q}_s}} \frac{\rho_L c^2}{\rho_S \Omega_L \Sigma} B_s^2(q_s).$$

To obtain formula (44) I have done the transition from discrete to continuous wave vectors:

$$\delta_{\mathbf{k}_{\parallel}, \mathbf{q}_{\alpha\parallel}} \longrightarrow \frac{(2\pi)^2}{\Sigma} \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{\alpha\parallel}).$$

The probability of creation of phonon with group velocity

$$\mathbf{c} \approx \frac{\omega_{\mathbf{k}}}{k} \frac{\mathbf{k}}{k} \quad (45)$$

near the surface element  $d\boldsymbol{\Sigma} \parallel \mathbf{e}_z$  ( $\mathbf{e}_z$  is the unit vector along  $z$ th direction) is given by

$$W^{(1\text{ph})} \hbar \omega_{\mathbf{k}} n(T_S) \frac{(\mathbf{c} \cdot d\boldsymbol{\Sigma})}{c} \approx W^{(1\text{ph})} \hbar \omega_{\mathbf{k}} n(T_S) d\Sigma, \quad (46)$$

where  $n(T_S) = [\exp(\hbar \omega_{\mathbf{q}_\alpha}/k_b T_S) - 1]^{-1}$  is the equilibrium distribution function of phonons in the solid,  $T_S$  is the temperature of the solid. Thus, the heat flux from solid to liquid helium per unit area can be written as follows

$$\begin{aligned} Q_{S \rightarrow L}^{(1\text{ph})} = (2\pi)^3 \Bigg\{ & \int \mathcal{A}_l \hbar \omega_{\mathbf{k}} n(T_S) \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{l\parallel}) \delta(\omega_{\mathbf{q}_l} - \omega_{\mathbf{k}}) \frac{\Omega_L d^3 k}{(2\pi)^3} \frac{\Omega_S d^3 q_l}{(2\pi)^3} \\ & + \int \mathcal{A}_t \hbar \omega_{\mathbf{k}} n(T_S) \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{t\parallel}) \delta(\omega_{\mathbf{q}_t} - \omega_{\mathbf{k}}) \frac{\Omega_L d^3 k}{(2\pi)^3} \frac{\Omega_S d^3 q_t}{(2\pi)^3} \\ & + \int \mathcal{A}_s \hbar \omega_{\mathbf{k}} n(T_S) \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{s\parallel}) \delta(\omega_{\mathbf{q}_s} - \omega_{\mathbf{k}}) \frac{\Omega_L d^3 k}{(2\pi)^3} \frac{\Sigma d^2 q_s}{(2\pi)^2} \Bigg\}. \end{aligned} \quad (47)$$

Delta functions of the type  $\delta(\mathbf{k}_{\parallel} - \mathbf{q}_{\alpha\parallel})$  take away integrations over  $\mathbf{k}_{\parallel}$ . Delta functions  $\delta(\omega_{\mathbf{q}_\alpha} - \omega_{\mathbf{k}})$  can be excluded by the integration over  $k_z$  (note,  $k_z \approx k$ ). After these straightforward calculations the integrals over the energies and over angles can be separated, so that the final result takes the form

$$Q_{S \rightarrow L}^{(1\text{ph})} = \frac{4\pi\hbar}{(2\pi)^3} \frac{\rho_L c}{\rho_S} \frac{F(\eta)}{s_t^3} \int_0^\infty \frac{\omega^3 d\omega}{\exp\left(\frac{\hbar\omega}{k_b T_S}\right) - 1} = \frac{\pi^2}{30} \frac{\rho_L c}{\hbar^3 \rho_S s_t^3} (k_b T_S)^4 F(\eta), \quad (48)$$

where  $F(\eta)$  is the function of  $\eta = s_l/s_t$  written down in Ref. 4. The formula (48) exactly coincides with the well known Khalatnikov result<sup>4</sup> derived initially in the framework of acoustic mismatch theory. Note, the total heat flux between the solid and liquid helium is given by the difference

$$J^{(1\text{ph})} = Q_{S \rightarrow L}^{(1\text{ph})} - Q_{L \rightarrow S}^{(1\text{ph})}, \quad (49)$$

where  $Q_{L \rightarrow S}^{(1\text{ph})}$  can be found in the similar way. Finally it is given by the same formula as (48) up to the replacement  $T_S \rightarrow T_L$ , where  $T_L$  is the temperature of liquid helium.

## V. ROTONS EMISSION

The novel techniques developed during last decade provide a possibility to produce and unambiguously detect by quantum evaporation processes both the  $R^{(-)}$  rotons ( $k < k_0$ ) and the  $R^{(+)}$  rotons ( $k > k_0$ ). However, it was realized that simple flat metal films produce easily measurable evaporation signals due to  $R^{(+)}$  rotons<sup>20</sup>, but no signal from  $R^{(-)}$  rotons has ever been identified from such sources. To produce  $R^{(-)}$  rotons two types of sources have been developed<sup>21,22</sup>, which are based on trapping of the emitted  $R^{(+)}$  rotons to enable  $R^{(-)}$  rotons to be created by interactions in nonequilibrium gas of excitations. Furthermore, Tucker and Wyatt reported<sup>22</sup> about creation of fast enough pulsed source of  $R^{(-)}$  rotons suitable for time of flight measurements.

Let us at first demonstrate the impossibility of direct emission of  $R^{(-)}$  rotons from a solid, and, secondly, to describe the process of  $R^{(+)}$  rotons emission.

### A. $R^{(-)}$ rotons

It is easy to see that  $R^{(-)}$  rotons and  $R^{(+)}$  rotons are not principally distinguished neither in the stress tensor (40) nor in the corresponding transfer Hamiltonian (41). Really the probability of  $R^{(-)}$  roton emission seems to be nonzero if the energy and parallel component of momentum are conserved. This is just the case in the emission of real particles, say quantum evaporation<sup>23</sup>. But quasiparticles in media are the *collective* excitations, which originated from local spatial perturbations. In this case the conservation laws at the surface should be obtained directly from the boundary conditions. In fact, the boundary conditions (6) and (7) just led us to the interaction Hamiltonian (8) we

have used. But the reversed conditions (which are clearly absurd) will necessarily give the same expression (8)! This means that for accurate use of this interaction Hamiltonian we must always keep in mind at least one of the boundary conditions, say (6) (the second condition (7) is automatically satisfied as long as we use the transfer Hamiltonian (8)). In original SBT theory<sup>13</sup> there were no boundary conditions at all. This is the reason why the problem of  $R^{(-)}$  rotons creation even did not appear in Ref. 13.

Let me now analyze the first boundary condition (6). In view of the expression (24) the normal component of displacement  $\delta r_z(x, y, 0)$  at the surface can be written as follows<sup>24</sup>

$$\delta r_z(x, y, 0) = \left( \frac{\hbar}{2\rho_L\Omega_L} \right)^{1/2} \sum_{\mathbf{k}} \omega_{\mathbf{k}}^{-1/2} \frac{k_z}{k} \left( \hat{a}_{\mathbf{k}} - \hat{a}_{-\mathbf{k}}^\dagger \right) e^{i\mathbf{k}_{\parallel}\mathbf{r}_{\parallel}}, \quad (50)$$

while retaining for simplicity one polarization, say longitudinal, eq. (25) at the interface can be rewritten in the analogous form:

$$u_z(x, y, 0) = \left( \frac{\hbar}{2\rho_S\Omega_S} \right)^{1/2} \sum_{\mathbf{q}} \omega_{\mathbf{q}}^{-1/2} \frac{q_z}{q} \left( \hat{b}_{\mathbf{q}} - \hat{b}_{-\mathbf{q}}^\dagger \right) e^{i\mathbf{q}_{\parallel}\mathbf{R}_{\parallel}}. \quad (51)$$

The right hand sides of eqs. (50) and (51) can be equal only if all their matrix elements in arbitrary basis are equal. Just this requirement leads to the conservation of the component of momentum parallel to the interface. But this can only occur if the coefficients at exponents in eqs. (50) and (51) have the same sign!

The detectable quasiparticle propagating into He II must have the positive  $z$ th component of group velocity. As is known momentum of  $R^{(-)}$  rotons is oppositely directed to their group velocity. This results in impossibility to satisfy the boundary condition (6) in view of eqs. (50) and (51).

### B. $R^{(+)}$ rotons emission

Let us now concentrate our attention in the processes of  $R^{(+)}$  rotons emission. In analogy with the previous section we can calculate the probability density function for single roton creation as

$$\begin{aligned} W^{(1R)} &= \frac{2\pi}{\Sigma\hbar^2} \left| \mathcal{M}_{\mathbf{q} \leftrightarrow \mathbf{k}}^{(1R)} \right|^2 \delta(\omega_{\mathbf{q}} - \omega_{\mathbf{k}}) \\ &= (2\pi)^3 \left\{ \mathcal{A}_l^{(1R)} \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{l\parallel}) + \mathcal{A}_t^{(1R)} \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{t\parallel}) + \mathcal{A}_s^{(1R)} \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{s\parallel}) \right\} \delta(\omega_{\mathbf{q}} - \omega_{\mathbf{k}}), \end{aligned} \quad (52)$$

where  $\mathbf{k}$  denotes now the roton wave vector,  $\omega_{\mathbf{k}}$  is its energy. The coefficients  $\mathcal{A}_l^{(1R)}$  are determined by the relations

$$\mathcal{A}_l^{(1R)} = \frac{\rho_L}{\rho_S\Omega_L\Omega_S} \frac{k^2}{4m^2} \frac{B_l^2(\theta)F^2(\mathbf{k})}{\omega_{\mathbf{k}}\omega_{\mathbf{q}_l}}, \quad \mathcal{A}_t^{(1R)} = \frac{\rho_L}{\rho_S\Omega_L\Omega_S} \frac{k^2}{4m^2} \frac{B_t^2(\theta)F^2(\mathbf{k})}{\omega_{\mathbf{k}}\omega_{\mathbf{q}_t}},$$

$$\mathcal{A}_s^{(1R)} = \frac{\rho_L}{\rho_S\Omega_L\Sigma} \frac{k^2}{4m^2} \frac{B_s^2(q_s)F^2(\mathbf{k})}{\omega_{\mathbf{k}}\omega_{\mathbf{q}_s}}.$$

The probability of creation of roton with energy  $\hbar\omega_{\mathbf{k}}$  and group velocity

$$\mathbf{v}_r = \frac{\partial\omega_{\mathbf{k}}}{\partial\mathbf{k}}$$

near the surface element  $d\Sigma \parallel \mathbf{e}_z$  is equal to

$$W^{(1R)} \hbar\omega_{\mathbf{k}} n(T_S) \frac{(\mathbf{v}_r \cdot d\Sigma)}{v_r} = W^{(1R)} \hbar\omega_{\mathbf{k}} n(T_S) \cos\theta d\Sigma, \quad (53)$$

where  $\cos\theta = \mathbf{v}_r \cdot \mathbf{e}_z$ . Therefore, the energy flux per unit area associated with single roton channel takes the form

$$\begin{aligned}
Q_{S \rightarrow L}^{(1R)} = (2\pi)^3 & \left\{ \int \mathcal{A}_i^{(1R)} \hbar \omega_{\mathbf{k}} \cos \theta n(T_S) \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{i\parallel}) \delta(\omega_{\mathbf{q}_i} - \omega_{\mathbf{k}}) \frac{\Omega_L d^3 k}{(2\pi)^3} \frac{\Omega_S d^3 q_i}{(2\pi)^3} \right. \\
& + \int \mathcal{A}_t^{(1R)} \hbar \omega_{\mathbf{k}} \cos \theta n(T_S) \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{t\parallel}) \delta(\omega_{\mathbf{q}_t} - \omega_{\mathbf{k}}) \frac{\Omega_L d^3 k}{(2\pi)^3} \frac{\Omega_S d^3 q_t}{(2\pi)^3} \\
& \left. + \int \mathcal{A}_s^{(1R)} \hbar \omega_{\mathbf{k}} \cos \theta n(T_S) \delta(\mathbf{k}_{\parallel} - \mathbf{q}_{s\parallel}) \delta(\omega_{\mathbf{q}_s} - \omega_{\mathbf{k}}) \frac{\Omega_L d^3 k}{(2\pi)^3} \frac{\Sigma d^2 q_s}{(2\pi)^2} \right\}. \quad (54)
\end{aligned}$$

In principle the partial integrations in eq. (54) can give the spectral and angular distributions of emitted rotons. Here I calculate the total energy yield from single roton channel. The account of all phonon polarizations in the solid can only give the multiplier similar to  $F(\eta)$  appeared in eq. (48). This multiplier is of the order of unity for majority of solids. The account of all polarizations can have an essential influence only on the angular distribution. So, to calculate the total energy flux let us restrict ourselves to one polarization, say longitudinal one. Further I drop all the polarization subscripts. Thus, the energy flux of interest looks

$$\begin{aligned}
Q_{S \rightarrow L}^{(1R)} = \frac{\hbar \rho_L}{4(2\pi)^3 m^2 \rho_S} \int \frac{k^2 \omega_{\mathbf{q}} \omega_{\mathbf{k}}^{-2} \cos \theta}{\exp\left(\frac{\hbar \omega_{\mathbf{q}}}{k_b T_S}\right) - 1} \left\{ \frac{m \omega_{\mathbf{k}}^2}{k^2} + \frac{\hbar^2 (k_z^2 - k^2)}{4m} \right\}^2 \\
\times \delta(\mathbf{q}_{\parallel} - \mathbf{k}_{\parallel}) \delta(\omega_{\mathbf{q}} - \omega_{\mathbf{k}}) n(T_S) d^3 k d^3 q. \quad (55)
\end{aligned}$$

After the straightforward, but quite cumbersome calculations eq. (55) can be reduced to the single integral. Let me write down here the spectral distribution for rotons emission:

$$\frac{dQ_{S \rightarrow L}^{(1R)}}{dk} = \frac{\rho_L \hbar^5}{120(2\pi)^2 m^4 s^7 \rho_S} \frac{k^2 \omega_{\mathbf{k}}^5}{\exp\left(\frac{\hbar \omega_{\mathbf{k}}}{k_b T_S}\right) - 1} \left\{ 1 - 10 \frac{m^2 s^2}{\hbar^2 k^2} + 30 \frac{m^4 s^4}{\hbar^4 k^4} \right\}, \quad (56)$$

To obtain the total energy flux via the single roton channel  $Q_{S \rightarrow L}^{(1R)}$ , the relation (56) should be integrated over  $k$  in the range  $\{k_0, k_{\max}\}$ , where  $k_{\max}$  is the maximum roton wave vector. Note again that the formula (56) gives the energy flux in the case  $T_L = 0$ . In general,  $Q_{L \rightarrow S}^{(1R)}$  can be calculated in the same way.

To make a numerical analysis of the obtained results we must choose the particular analytic expression for roton spectrum in He II. Precise numerical calculations require of course the analytic expression most closed to a reality. But to draw the general picture I will use here the classical Landau expression<sup>1</sup>

$$\hbar \omega_{\mathbf{k}} = \Delta + \frac{\hbar^2 (k - k_0)^2}{2\mu} \quad (57)$$

where  $\Delta \approx 8.6$  K,  $k_0 \approx 1.9 \cdot 10^8$  cm<sup>-1</sup>,  $\mu \approx 0.13 m_{\text{He}^4} \approx 0.868 \cdot 10^{-24}$  g.

Figure 2 presents the spectral distribution of roton emission calculated from (56) for gold. Figure clearly shows that the maximum at  $k > k_0$  appears when temperature of the solid exceed 2 K. Integrating eq. (56) over  $R^{(+)}$  rotons momenta we get the total energy flux. The ratio of this flux associated with the single roton processes and that corresponding to the single phonon processes given by eq. (48) is shown in Figure 2 as a function of temperature of the solid (again gold). Previously SBT<sup>13</sup> had also estimated the contribution of single roton emission processes into the total energy transmission through the solid – liquid helium interface. They, however, considered the case of small differences between  $T_S$  and  $T_L$ . Corresponding formulae can be easily obtained from eqs. (56) and (48) by differentiation with respect to a temperature and subsequent integration over roton and phonon momenta, respectively. The corresponding ratio is shown in Figure 3 together with SBT result. The estimation by SBT<sup>13</sup> gives the ratio approximately three times larger for  $T_S = 2$  K. This difference is caused by the approximations done in Ref. 13. Particularly the authors had taken  $k_z \approx k$ , which is far from reality for rotons. Additionally they included  $R^{(-)}$  rotons in their calculations, which is incorrect as I show in the previous subsection. But one can see from Fig. 3 that their rough estimation nevertheless show right qualitative behaviour of the energy flux due to roton emission.

## VI. CONCLUSIONS

I studied the emission of elementary excitations into liquid helium II from a heated solid. For this process I developed the new microscopic quantum theory generalizing the previous theory<sup>13</sup>. To describe the coupling between

the solid and He II the correct expression for the interface Hamiltonian (8) has been derived. This expression follows from the general principles of quantum mechanics with account of boundary conditions at solid–liquid interface (6) and (7). Following the ideas of Ref. 13 and using the microscopic description of superfluid helium<sup>15</sup> the transfer Hamiltonian (8) has been expressed in terms of creation and annihilation operators of elementary excitations in the solid and helium II. The transfer Hamiltonians are explicitly written down for i) single phonon emission (34), ii) two phonons emission (36), and iii) single roton emission (41). The heat flux through the interface associated with the single phonon channel is calculated by the use of the derived microscopic expression (34). The obtained formula (48) exactly coincides with the corresponding result obtained by Khalatnikov<sup>4</sup> using the acoustic-mismatch approach. The heat flux associated with the single  $R^{(+)}$  roton channel (see formulas (55) and (56)) has been calculated as well. It is shown that the boundary conditions (6) and (7) forbid the emission of  $R^{(-)}$  rotons by a solid. This latter result is in accordance with experimental observations.

## ACKNOWLEDGMENTS

I would like to acknowledge the numerous stimulating discussions on the subject with Prof. I. N. Adamenko and Dr. K. É. Nemchenko. I wish to express also my deep gratitude to Prof. S. Yamasaki and Prof. M. Hirooka who acquainted me with the papers cited in Ref. 15.

- 
- <sup>1</sup> L. D. Landau, J. Phys. USSR **5**, 71 (1941); **11**, 91 (1947)  
<sup>2</sup> R. P. Feynman and M. Cohen, Phys. Rev. **102**, 1183 (1956)  
<sup>3</sup> P. L. Kapitza, J. Phys. USSR, **4**, 181 (1941)  
<sup>4</sup> I. M. Khalatnikov, Zh. Eksp. Teor. Fiz. **22**, 687 (1952); I. M. Khalatnikov, *Introduction to the Theory of Superfluidity*, (Benjamin, New York, 1965); see also the corrected and augmented monograph in Russian: I. M. Khalatnikov, *Theory of Superfluidity*, (Nauka, Moscow, 1971)  
<sup>5</sup> W. A. Little, Can. J. Phys. **37**, 334 (1959)  
<sup>6</sup> L. J. Challis, J. Phys. C: Solid State Phys. **7**, 481 (1974)  
<sup>7</sup> K. N. Zinov'eva, Fiz. Nizk. Temp. **23**, 485 (1997) [English translation: Low. Temp. Phys. **23**, 355 (1997)]  
<sup>8</sup> M. A. H. Tucker, A. F. G. Wyatt, I. N. Adamenko, A. V. Zhukov, and K. É. Nemchenko, Fiz. Nizk. Temp. **25**, 657 (1999) [English translation: Low. Temp. Phys. **25**, 488 (1999)]; M. A. H. Tucker, A. F. G. Wyatt, I. N. Adamenko, K. É. Nemchenko, and A. V. Zhukov, Phys. Rev. B **62**, 3029 (2000)  
<sup>9</sup> R. A. Sherlock, A. F. G. Wyatt, N. G. Mills, and N. A. Lockerbie, Phys. Rev. Lett. **29**, 1299 (1972)  
<sup>10</sup> R. A. Sherlock, N. G. Mills, and A. F. G. Wyatt, J. Phys. C: Solid State Phys. **8**, 300 (1975)  
<sup>11</sup> L. J. Challis, K. Dransfeld, and J. Wilks, Proc. Roy. Soc. A **260**, 31 (1961); A. F. Andreev, Zh. Eksp. Teor. Phys. **43**, 1535 (1962) [English translation: Sov. Phys. JETP **16**, 1084 (1963)]; I. N. Adamenko and I. M. Fuks, Zh. Eksp. Teor. Phys. **59**, 2071 (1970) [English translation: Sov. Phys. JETP **32**, 1123 (1971)]; I. M. Khalatnikov and I. N. Adamenko, Zh. Eksp. Teor. Phys. **63**, 745 (1972) [English translation: Sov. Phys. JETP **36**, 391 (1973)]; R. E. Peterson and A. C. Anderson, Phys. Lett. A **40**, 317 (1972); Solid State Commun. **10**, 891 (1972); H. Haug and K. Weiss, Phys. Lett. A **40**, 19 (1972); M. Vuorio, J. Low Temp. Phys. **10**, 781 (1973)  
<sup>12</sup> G. A. Toombs and L. J. Challis, J. Phys. C: Solid State Phys. **4**, 1085 (1971); M. J. Rice, Phys. Rev. Lett. **27**, 92 (1971); F. W. Sheard, G. A. Toombs, and L. J. Challis, Phys. Rev. Lett. **27**, 1117 (1971); M. J. Rice, and G. A. Toombs, Phys. Rev. A **5**, 2259 (1972); G. A. Toombs, F. W. Sheard, and M. J. Rice, Solid State Commun. **10**, 601 (1972)  
<sup>13</sup> F. W. Sheard, R. M. Bowley, and G. A. Toombs, Phys. Rev. A **8**, 3135 (1973)  
<sup>14</sup> M. Brown and A. F. G. Wyatt, J. Phys.: Cond. Matt. **2**, 5025 (1990); A. C. Forbes and A. F. G. Wyatt, Phys. Rev. Lett. **64**, 1393 (1990); see also Refs. 20–22.  
<sup>15</sup> S. Sunakawa, S. Yamasaki, and T. Kebukawa, Prog. Theor. Phys. **41**, 919 (1969); see also the generalization of this earlier theory in the subsequent series of papers S. Yamasaki, T. Kebukawa, and S. Sunakawa, Prog. Theor. Phys. **53**, 1243 (1975); Prog. Theor. Phys. **54**, 348 (1975); Prog. Theor. Phys. **56**, 415 (1976); Prog. Theor. Phys. **61**, 996 (1979); Prog. Theor. Phys. **61**, 1012 (1979); S. Yamasaki and M. Hirooka, Prog. Theor. Phys. **64**, 1248 (1980); M. Hirooka and S. Yamasaki, Prog. Theor. Phys. **64**, 1521 (1980); S. Yamasaki and M. Hirooka, Prog. Theor. Phys. **66**, 1200 (1981); Prog. Theor. Phys. **67**, 20 (1982); M. Hirooka and S. Yamasaki, Prog. Theor. Phys. **68**, 1050 (1982); Prog. Theor. Phys. **68**, 1482 (1982); S. Yamasaki, Prog. Theor. Phys. **72**, 441 (1984); Prog. Theor. Phys. **73**, 598 (1985); Prog. Theor. Phys. **73**, 1320 (1985)  
<sup>16</sup> B. N. Esel'son, V. N. Grigor'ev, V. G. Ivantsov, É. Ya. Rudavskii, D. G. Sanikidze, and I. A. Serbin, *Mixtures of Quantum Liquids  $^3\text{He}-^4\text{He}$* , (Nauka, Moscow, 1973)

<sup>17</sup> L. P. Pitaevskii, Zh. Eksp. Teor. Fiz. **31**, 536 (1956)

<sup>18</sup> The expressions (23) and (24) are different in form from those in Refs. 13 and 15, because of using another notations. Such a method of presentation is closer to the ordinary quantization of liquid helium hydrodynamic Hamiltonian (see the first paper in Ref. 8).

<sup>19</sup> I. M. Khalatnikov and Yu. A. Matveev, J. Low Temp. Phys. **51**, 99 (1983)

<sup>20</sup> F. R. Hope, M. J. Baird, and A. F. G. Wyatt, Phys. Rev. Lett. **52**, 1528 (1984)

<sup>21</sup> H. Baddar, D. O. Edwards, T. M. Levin, and M. S. Pettersen, Physica B **194-196**, 513 (1994)

<sup>22</sup> M. A. H. Tucker and A. F. G. Wyatt, J. Low Temp. Phys. **110**, 455 (1998); **113**, 615 (1998); Science **283**, 1150 (1999)

<sup>23</sup> M. J. Baird, F. R. Hope, and A. F. G. Wyatt, Nature **304**, 325 (1983); A. F. G. Wyatt, Nature **391**, 56 (1998); F. Dalfovo, A. Fracchetti, A. Lastri, L. Pitaevskii, and S. Stringari, J. Low Temp. Phys. **104**, 367 (1996); F. Dalfovo, M. Guilleumas, A. Lastri, L. Pitaevskii, and S. Stringari, J. Phys.: Condens. Matter **9**, L369 (1997); C. D. H. Williams, J. Low Temp. Phys. **113**, 11 (1998); J. Low Temp. Phys. **113**, 627 (1998)

<sup>24</sup> Formula (50) becomes obvious if we rewrite collective variable (24) in Heisenberg representation simply multiplying  $\hat{a}_{\mathbf{k}}$  by  $\exp(-i\omega_{\mathbf{k}}t)$  and  $\hat{a}_{\mathbf{k}}^\dagger$  by  $\exp(i\omega_{\mathbf{k}}t)$ .

FIG. 1. Geometry of the problem

FIG. 2. Spectral distribution of roton emission into He II from golden heater calculated by eq. (56)

FIG. 3. Ratio of energy flux associated with the single roton channel and that corresponding to the single phonon (Khalatnikov) channel in case  $T_L = 0$

FIG. 4. Ratio of energy flux associated with the single roton channel and that corresponding to the single phonon (Khalatnikov) channel in case  $T_L \sim T_S$ . Corresponding result of SBT<sup>13</sup> is presented by the broken curve







